

# VIANT PULVER TIL INFUSJONSVÆSKE, OPPLØSNING

## VITAMINPREPARAT TIL PARENTERALT BRUK – VANNLØSELIGE- OG FETTLØSELIGE VITAMINER B05X C-

Vitamin	Anbefalinger	Viant
Tiamin (vitamin B <sub>1</sub> )	6 mg	6 mg
Riboflavin (vitamin B <sub>2</sub> )	3,6 mg	3,6 mg
Nikotinamid (vitamin B <sub>3</sub> )	40 mg	40 mg
Folsyre (vitamin B <sub>9</sub> )	600 µg	600 µg
Pantotensyre (vitamin B <sub>5</sub> )	15 mg	15 mg
Pyridoksin (vitamin B <sub>6</sub> )	6 mg	6 mg
Cyanokobalamin (vitamin B <sub>12</sub> )	5 µg	5 µg
Biotin (vitamin B <sub>7</sub> )	60 µg	60 µg
Askorbinsyre (vitamin C)	200 mg	200 mg
Retinol (vitamin A)	1 mg/3300 IU	1 mg/3300 IU
Kolekalsiferol (vitamin D)	5 µg	5 µg
Tokoferol (vitamin E)	10 mg/10 IU	10 mg/10IU
Fytomenadion (vitamin K <sub>1</sub> )	150 µg	150 µg

### Referanser:

1. ESPEN Guidelines on Parenteral Nutrition: Intensive care; P. Singer, M. M. Berger, G. V. d. Berghe, G. Biolo, P. Calder, A. Forbes, R. Griffiths, G. Kreyman, X. Lerverve, C. Pichard; *Clinical Nutrition* 28 (2009) 387–400.
2. ESPEN Guidelines on Parenteral Nutrition: Surgery; M. Braga, O. Ljungqvist, P. Soeters, K. Fearon, A. Weimann, F. Bozzetti; *Clinical Nutrition* 28 (2009) 378–386.
3. Food and Drug Administration (FDA). parenteral multivitamin products; drugs for human use; drug efficacy study implementation; amendment. *Federal Register* 2000;65:21200–1.
4. Micronutrient Supplementation in Adult Nutrition Therapy: Practical Considerations, Krishnan Sriram and Vassyl A. Lonchyna, *JPEN J Parenter Enteral Nutr* 2009 33: 548 originally published online 19 May 2009, DOI: 10.1177/0148607108328470.